

Preliminary analysis of the impact of airport closures on local air quality

We have received many enquiries asking what the effects of the closure of UK airspace has had on air quality surrounding major airports. In response we have made an initial analysis of NO_x and NO₂ concentrations surrounding Gatwick and Heathrow airports during the first three days of closure – Thursday 15th to Saturday 17th April 2010. This period was chosen due to the stable weather conditions with light north easterly winds, allowing a cross-sectional analysis upwind and downwind of the airports.

This period of unprecedented closure during unexceptional weather conditions has allowed us to demonstrate that the airports have a clear measurable effect on NO₂ concentrations and that this effect disappeared entirely during the period of closure, leading to a temporary but significant fall in pollutant concentrations adjacent to the airport perimeters.

Concentrations recorded by a monitoring site arise from a mix of pollution sources both local and distant affecting that specific location. In order to separate and quantify the affects of a particular local source, such as emissions from an airport, a control dataset is required. In this case pairs of monitoring sites were used located either side of each airport. The upwind monitors provided the control data. By subtracting hourly mean concentrations recorded by the upwind site from those recorded by the downwind site, an estimation of emissions from the airport could be made – termed ‘airport’ concentrations.

Gatwick Airport

Reigate and Banstead Borough Council operate three continuous monitoring sites close to the perimeter of Gatwick airport, two of which were well placed to make such a cross-sectional assessment. Poles Lane (‘RG3’) is situated in a suburban location approximately 0.5 km to the south west of the runway. This site provided downwind concentrations during north easterly winds. Horley (‘RG1’) is in a suburban location approximately 2.5 km to the north east of the runway and therefore provided upwind concentrations. Both sites continuously monitor NO_x and have been in operation since 2005 and 2000 respectively.

In order to assess the impact of the airport closure on NO_x and NO₂ concentrations, hourly mean ‘airport’ NO_x and NO₂ concentrations at RG3 were calculated from April 2005 up to the date of closure (15th April 2010 12:00) using the method described above. This process was then repeated for the closure days (15th April 2010 12:00 to 19th April 2010 00:00). Local wind data were taken from the RG1 and RG3 monitoring sites. This dataset was then filtered to only include data recorded during wind directions between 0 and 90 degrees when emissions arising within the airport had been shown to have an impact.

Figure 1 shows filtered daily diurnal mean ‘airport’ NO_x concentrations for the ‘open’ and ‘closed’ periods. The yellow shaded areas indicate hours where filtered data were available during the closure period, i.e., winds were from the north or north east. This chart shows that during normal operating conditions ‘airport’ NO_x concentrations increase during the day peaking at around 22 to 25 µg m⁻³ on average. Error bars show 95% confidence intervals, i.e., concentrations over the five year period fell between these error bars approximately 95% of the time. During the period of closure

(15th to 17th April 2010) mean 'airport' NO_x concentrations were zero most of the time, with a short peak of 4 µg m⁻³ on Saturday at 10am. All concentrations were well below the lower 95% confidence interval indicating that the difference from 'normal' operation was statistically significant.

The analysis was repeated for NO₂, with similar results. 'Airport' NO₂ concentrations dropped from a mean of 8 µg m⁻³ (peak of 13 µg m⁻³) during normal operation (06:00 to 22:00) to zero during most hours of closure (maximum of 1 µg m⁻³ on Friday at 01:00). The annual mean NO₂ concentration measured at RG3, to the south west of the airport, during 2009 would decrease from 18 µg m⁻³ to approximately 16 µg m⁻³ in the absence of airport emissions. The impact of the airport is likely to be greater in the populated areas to the north east of the airport (Horley) due to prevailing winds from the south west.




Figure 1: Comparison of diurnal variation in filtered 'airport' NO_x concentrations during normal operation and airport closure.

Heathrow Airport

A similar analysis was carried out using a NE/SW cross-section pair of monitoring sites surrounding Heathrow airport. The downwind site, Oaks Road ('LH7') is located approximately 0.6 km south of Heathrow's southern runway. Due to data not being available from closer sites to the north east of the airport, a site in Southall ('EA7') approximately 4.5 km from the airport was used as the upwind site. The data periods and filtered wind directions were the same as the Gatwick analysis. Due to slightly different wind patterns and an earlier change in wind direction, the analysis could only be carried out on Thursday and Friday 15th and 16th April.

During normal operating conditions the LH7 site recorded mean 'airport' NO_x concentrations of 64 µg m⁻³ (06:00 to 22:00), far higher than those recorded at the Gatwick site. This dropped to 10 µg m⁻³ during the closure period. 'Airport' NO₂ concentrations, which were also higher than those at Gatwick, dropped from 27 µg m⁻³ to 8 µg m⁻³. In the absence of airport emissions, the annual mean

NO₂ concentration recorded at LH7 during 2009 would decrease from as 33 µg m⁻³ to approximately 30 µg m⁻³. As with Gatwick, the impact of airport emissions is likely be greater to the north east of the airport over a full year.

This exceptional closure has allowed us to demonstrate the impacts of airport emissions on their immediate neighbourhood. The evidence from this preliminary analysis can be extended to quantify the impacts of Gatwick and Heathrow airport on their neighbourhoods during normal operation.

This preliminary study did not consider the impact of decreased traffic flows on airport feeder roads. Decreased flows are likely to have a significant effect on concentrations of vehicle-related pollutants close to such roads. Unfortunately, we do not have sufficient traffic data to carry out this analysis at this time.

Ben Barratt and Gary Fuller
King's College London Environmental Research Group